


Seattle and King County Monthly Syphilis Report: August 2022

Recent data are incomplete*

Syphilis Cases by Stage, King County September 2020-August 2022




*Note: All data are preliminary and data for past months may change due to delays in reporting from labs and providers. This report includes cases reported through September 16, 2022.


MSM = men who have sex with men
MSW = men who have sex with women only

Early syphilis includes primary, secondary, and early non-primary, non-secondary stages.

Early syphilis cases among cisgender men who have sex with men (MSM), King County September 2020-July 2022**



Early syphilis cases among cisgender men who have sex with women (MSW) and cisgender women, King County September 2020-July 2022**




**Due to incomplete reporting cases for the most recent month are not shown by gender and gender of sex partners.

Seattle and King County Monthly Syphilis Report: August 2022

Syphilis (all stages) cases by gender and gender of sex partners, January-August 2022

| | |
|---|-----|
| Cisgender MSM | 522 |
| Cisgender MSW | 203 |
| Cisgender Women | 213 |
| Cisgender Men, Unknown Gender of Sex Partners | 86 |
| Transgender, non-binary, or genderqueer | 31 |

Pregnant and congenital syphilis cases, King County September 2020-August 2022



25 syphilis cases were **pregnant** at the time of diagnosis and **8 congenital** cases have been diagnosed in 2022.

Syphilis (all stages) cases interviewed for partner services among cisgender men who have sex with men (MSM), cisgender men who have sex with women (MSW), cisgender women, Seattle and South King County September 2020-August 2022*

| Characteristics | Region | | | | | | | |
|-------------------------------------|---------|------|---------------|------|-------------------|------|---------------|------|
| | Seattle | | | | South King County | | | |
| | MSM | | MSW and Women | | MSM | | MSW and Women | |
| | N | % | N | % | N | % | N | % |
| Total Interviewed | 165 | 100% | 193 | 100% | 91 | 100% | 233 | 100% |
| Homeless (past 3 months) | 11 | 7% | 78 | 40% | 9 | 10% | 51 | 22% |
| Injection Drug Use (past year) | 7 | 4% | 21 | 11% | 1 | 1% | 10 | 4% |
| Methamphetamine Use (past year) | 29 | 18% | 61 | 32% | 12 | 13% | 52 | 22% |
| Exchange Sex (past year) | 5 | 3% | 21 | 11% | 4 | 4% | 9 | 4% |
| No Risk Identified** | 133 | 81% | 89 | 46% | 73 | 80% | 155 | 67% |
| Race/ethnicity | | | | | | | | |
| American Indian/Alaska Native | 2 | 1% | 5 | 3% | 0 | 0% | 3 | 1% |
| Asian | 9 | 5% | 11 | 6% | 5 | 5% | 9 | 4% |
| Black | 19 | 12% | 60 | 31% | 15 | 16% | 65 | 28% |
| Hispanic/Latinx | 36 | 22% | 33 | 17% | 26 | 29% | 58 | 25% |
| Native Hawaiian or Pacific Islander | 3 | 2% | 3 | 2% | 2 | 2% | 6 | 3% |
| White | 81 | 49% | 58 | 30% | 36 | 40% | 66 | 28% |
| Multiracial | 4 | 2% | 9 | 5% | 2 | 2% | 11 | 5% |
| Other/Unknown | 11 | 7% | 14 | 7% | 5 | 5% | 15 | 6% |
| Age | | | | | | | | |
| 15-19 | 3 | 2% | 3 | 2% | 2 | 2% | 9 | 4% |
| 20-24 | 22 | 13% | 21 | 11% | 15 | 16% | 35 | 15% |
| 25-29 | 36 | 22% | 29 | 15% | 13 | 14% | 45 | 19% |
| 30-34 | 29 | 18% | 31 | 16% | 28 | 31% | 50 | 21% |
| 35-44 | 40 | 24% | 58 | 30% | 16 | 18% | 54 | 23% |
| 45-54 | 18 | 11% | 29 | 15% | 8 | 9% | 25 | 11% |
| >=55 | 17 | 10% | 22 | 11% | 9 | 10% | 15 | 6% |

*Note: All data are preliminary and data for past months may change due to delays in reporting from labs and providers. This report includes cases reported through September 16, 2022. The table above is limited to cases who were interviewed by Public Health staff.

**No reported homelessness, injection drug use, methamphetamine use, or exchange sex.