



Medicine for Latent TB: 3HP

You will take your medicine once per week for 12 weeks.

You will take your weekly pills together at the same time, on the same day of the week.

Tell your other doctors that you are taking medicine for latent TB (LTBI).



You have latent TB, a kind of TB infection. You do not have TB disease and cannot spread TB to other people.

This medicine will help prevent active TB disease.



Call your nurse if you have questions or side effects:


Medicine	Number of pills	Day of the week
Isoniazid		M T W Th F Sat Sun
Rifapentine		M T W Th F Sat Sun
Pyridoxine (vit B6)		M T W Th F Sat Sun



STOP taking your medicine right away and CALL your TB nurse if you have any of the problems below:

- Little or no appetite for food
- An upset stomach or stomach cramps
- Nausea or vomiting
- Cola-colored urine or light stools
- Yellowing skin or eyes
- Easy bruising or bleeding
- Rash or itching
- Severe weakness or tiredness
- Fever
- Head or body aches
- Dizziness
- Tingling or numbness in your hands or feet

NOTE: It is OK if your urine, saliva, or tears turn orange. Soft contact lenses may become stained.



Tips for taking 3HP

- Set an alarm to remind you to take your medicine
- Ask a family or friend to remind you
- Talk to your doctor if you drink alcohol, alcohol may cause side effects
- Tell your doctor about other medicine you take
- Ask your nurse if you have questions